

MSUMB 97-09, December 1997

Hopf algebra structure of $\mathrm{Gr}_q(1|1)$ related to $\mathrm{GL}_q(1|1)$

Salih Celik ^{1 2}

Mimar Sinan University, Department of Mathematics,
80690 Besiktas, Istanbul, TURKEY.

Abstract

We show that the algebra of functions on the Grassmann supergroup $\mathrm{Gr}_q(1|1)$ has a (graded) Hopf algebra structure related to $\mathrm{GL}_q(1|1)$.

¹E-mail: scelik@fened.msu.edu.tr

²E-mail: sacelik@yildiz.edu.tr

In the past few years, quantum groups¹ and q -deformed universal enveloping algebras² have been intensively studied both by mathematicians and mathematical physicists. From a mathematical point of view, these algebraic structures are just special classes of noncommutative Hopf algebras.

The algebraic structure underlying quantum groups extends the theory of the supergroups.³ The simplest quantum supergroup is $GL_q(1|1)$, i.e. the deformation of the supergroup of 2x2 matrices with two bosonic (even) and two fermionic (odd) matrix entries.

The aim of the present work is to construct the (graded) Hopf algebra structure of the Grassmann supergroup $Gr_q(1|1)$, the superdual of $GL_q(1|1)$, which was introduced in Ref. 4. Before discussing the (graded) Hopf algebra structure of $Gr_q(1|1)$, let us first give some notations and useful formulas about the quantum Grassmann supergroup $Gr_q(1|1)$.

A Grassmann supermatrix $\hat{T}$ which is an element of $Gr(1|1)$, is of the form

$$\hat{T} = \begin{pmatrix} \alpha & b \\ c & \delta \end{pmatrix}$$

with two odd (greek letters) and two even (latin letters) matrix elements. The symbol *hat* is used to distinguish $\hat{T}$ from an element T of $GL_q(1|1)$.

The q -deformation of the Grassman supergroup $Gr(1|1)$ as a quantum matrix supergroup $Gr_q(1|1)$ is generated by α, b, c, δ with the relations⁴ :

$$\begin{aligned} \alpha b &= q^{-1} b \alpha, & \alpha c &= q^{-1} c \alpha, \\ \delta b &= q^{-1} b \delta, & \delta c &= q^{-1} c \delta, \\ \alpha \delta + \delta \alpha &= 0, & \alpha^2 &= 0 = \delta^2, \\ bc &= cb + (q - q^{-1}) \delta \alpha \end{aligned} \tag{1}$$

where q is a non-zero complex number and $q^2 \neq 1$. The associative algebra (1) is equivalent to equation⁵

$$R^1 \hat{T}_1 \hat{T}_2 = -\hat{T}_2 \hat{T}_1 R^2 \tag{2}$$

where

$$R^1 = \begin{pmatrix} q & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & q - q^{-1} & -1 & 0 \\ 0 & 0 & 0 & q^{-1} \end{pmatrix}, \quad R^2 = \begin{pmatrix} q^{-1} & 0 & 0 & 0 \\ 0 & -1 & q^{-1} - q & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & q \end{pmatrix}, \tag{3}$$

are both the solutions of the quantum (graded) Yang-Baxter equation. Here, we used the tensoring convention

$$\begin{aligned}(\hat{T}_1)_{kl}^{ij} &= (\hat{T} \otimes I)_{kl}^{ij} = \hat{T}_k^i \delta_l^j, \\ (\hat{T}_2)_{kl}^{ij} &= (I \otimes \hat{T})_{kl}^{ij} = (-1)^{i(j+l)} \hat{T}_l^j \delta_k^i.\end{aligned}\tag{4}$$

The central element of the algebra (1) is⁴

$$\hat{D}_q = bc^{-1} - \alpha c^{-1} \delta c^{-1} = c^{-1}b - c^{-1}\alpha c^{-1}\delta.\tag{5}$$

We now denote the algebra generated by the elements α, b, c, δ with the relations (1) by $\hat{\mathcal{A}}$. We want to make the algebra $\hat{\mathcal{A}}$ into a (graded) Hopf algebra related to the quantum supergroup $\text{GL}_q(1|1)$. Because of this, we state briefly some properties of the quantum supergroup $\text{GL}_q(1|1)$ we are going to need in this work.

The quantum supergroup $\text{GL}_q(1|1)$ is generated by four generators a, β, γ, d and the q -commutation relations³

$$\begin{aligned}a\beta &= q\beta a, & d\beta &= q\beta d, \\ a\gamma &= q\gamma a, & d\gamma &= q\gamma d, \\ \beta\gamma + \gamma\beta &= 0, & \beta^2 &= 0 = \gamma^2, \\ ad &= da + (q - q^{-1})\gamma\beta.\end{aligned}\tag{6}$$

The generators satisfying the relations (6) generate the algebra called the algebra of functions on the quantum supergroup $\text{GL}_q(1|1)$ and we shall denote it by $\mathcal{A}$. We know that the algebra $\mathcal{A}$ is a (graded) Hopf algebra whose structure we now discuss. We represent the set of generators a, β, γ, d in the form of a matrix

$$T = \begin{pmatrix} a & \beta \\ \gamma & d \end{pmatrix}.$$

Then the relations (6) are equivalent to equation

$$RT_1T_2 = T_2T_1R\tag{7}$$

where

$$R = \begin{pmatrix} q & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & q - q^{-1} & 1 & 0 \\ 0 & 0 & 0 & q^{-1} \end{pmatrix}.\tag{8}$$

The superinverse of T is given by³

$$T^{-1} = \begin{pmatrix} A & \Omega \\ \Gamma & D \end{pmatrix} = \begin{pmatrix} a^{-1} + a^{-1}\beta d^{-1}\gamma a^{-1} & -a^{-1}\beta d^{-1} \\ -d^{-1}\gamma a^{-1} & d^{-1} + d^{-1}\gamma a^{-1}\beta d^{-1} \end{pmatrix},\tag{9}$$

and the superdeterminant is

$$D_q = ad^{-1} - \beta d^{-1} \gamma. \quad (10)$$

It is easy to verify that D_q commutes with all matrix elements of T . Note that the matrix elements of T with those of T^{-1} satisfy the following relations

$$\begin{aligned} aA &= q^2 Aa + 1 - q^2, & dA &= Ad, \\ aD &= Da, & dD &= q^2 Dd + 1 - q^2, \\ a\Omega &= q\Omega a, & d\Omega &= q\Omega d, \\ a\Gamma &= q\Gamma a, & d\Gamma &= q\Gamma d, \\ \beta A &= qA\beta, & \gamma A &= qA\gamma, \\ \beta D &= qD\beta, & \gamma D &= qD\gamma, \\ \beta\Omega &= \Omega\beta, & \gamma\Omega &= -q^2\Omega\gamma, \\ \beta\Gamma &= -q^2\Gamma\beta, & \gamma\Gamma &= \Gamma\gamma. \end{aligned} \quad (11)$$

The usual coproduct is given by

$$\Delta : \mathcal{A} \longrightarrow \mathcal{A} \otimes \mathcal{A}, \quad \Delta(t_j^i) = t_k^i \otimes t_j^k \quad (12)$$

where summation over repeated indices is understood. One can rewrite the last formula in the following nice and elegant form

$$\Delta(T) = T \dot{\otimes} T \quad (13)$$

where $\otimes$ stands for the usual tensor product and the dot refers to the summation over repeated indices and reminds us about the usual matrix multiplication. The counit is given by

$$\varepsilon : \mathcal{A} \longrightarrow \mathcal{C}, \quad \varepsilon(t_j^i) = \delta_j^i. \quad (14)$$

The coinverse (antipode) is given by

$$S : \mathcal{A} \longrightarrow \mathcal{A}, \quad S(T) = T^{-1}. \quad (15)$$

It is not difficult to verify the following properties of the co-structures:

$$(\Delta \otimes \text{id}) \circ \Delta = (\text{id} \otimes \Delta) \circ \Delta, \quad (16)$$

$$\mu \circ (\varepsilon \otimes \text{id}) \circ \Delta = \mu' \circ (\text{id} \otimes \varepsilon) \circ \Delta, \quad (17)$$

$$m \circ (S \otimes \text{id}) \circ \Delta = \varepsilon = m \circ (\text{id} \otimes S) \circ \Delta, \quad (18)$$

where id denotes the identity mapping,

$$\mu : \mathcal{C} \otimes \mathcal{A} \longrightarrow \mathcal{A}, \quad \mu' : \mathcal{A} \otimes \mathcal{C} \longrightarrow \mathcal{A}$$

are the canonical isomorphisms, defined by

$$\mu(k \otimes a) = ka = \mu'(a \otimes k), \quad \forall a \in \mathcal{A}, \quad \forall k \in \mathcal{C} \quad (20)$$

and m is the multiplication map

$$m : \mathcal{A} \otimes \mathcal{A} \longrightarrow \mathcal{A}, \quad m(a \otimes b) = ab. \quad (21)$$

The multiplication in $\mathcal{A} \otimes \mathcal{A}$ follows the rule

$$(A \otimes B)(C \otimes D) = (-1)^{p(B)p(C)} AC \otimes BC \quad (22)$$

where $p(X)$ is the z_2 -grade of X , i.e. $p(X) = 0$ for even variables and $p(X) = 1$ for odd variables.

Since the (graded) Hopf algebra structure of $\text{Gr}_q(1|1)$ is related to those of $\text{GL}_q(1|1)$ it is necessary to obtain the commutation relations of the generators of $\hat{\mathcal{A}}$ with those of $\mathcal{A}$. We define the (mixed) commutation relations between the generators of $\hat{\mathcal{A}}$ and $\mathcal{A}$ as follows:

$$R\hat{T}_1T_2 = (-1)^{p(T_2)}T_2\hat{T}_1R', \quad R' = R - (q - q^{-1})P \quad (23)$$

where P is the superpermutation matrix. The equation (23) gives the mixed relations

$$\begin{aligned} a\alpha &= q^2\alpha a, & \beta\alpha &= -q\alpha\beta, \\ ab &= qba + (q^2 - 1)\alpha\beta, & \beta b &= b\beta, \\ ac &= qca + (q^2 - 1)\alpha\gamma, & \beta c &= c\beta + (q - q^{-1})\alpha d, \\ a\delta &= \delta a + (q - q^{-1})(\beta c - b\gamma), & \beta\delta &= -q^{-1}\delta\beta + (1 - q^{-2})bd, \\ d\alpha &= \alpha d, & \gamma\alpha &= -q\alpha\gamma, \\ db &= q^{-1}bd, & \gamma b &= b\gamma - (q - q^{-1})\alpha d, \\ dc &= q^{-1}cd, & \gamma c &= c\gamma, \\ d\delta &= q^{-2}\delta d, & \gamma\delta &= -q^{-1}\delta\gamma + (1 - q^{-2})cd. \end{aligned} \quad (24)$$

Using these relations, it is easy to verify that $\hat{D}_q$, which is given by (5), is still a central element, i.e. $\hat{D}_q$ also commutes with the generators of $\mathcal{A}$.

After some algebra the commutation relations between the matrix elements of $\hat{T}$ with T^{-1} are obtained to be

$$\alpha A = q^2 A \alpha, \quad \delta A = A \delta,$$

$$\begin{aligned}
\alpha D &= D\alpha, & \delta D &= q^{-2}D\delta + (q - q^{-1})^2 A\alpha + (q^{-2} - 1)(\Omega c - \Gamma b), \\
\alpha \Omega &= -q\Omega\alpha, & \delta \Omega &= -q^{-1}\Omega\delta + (q^{-1} - q)Ab, \\
\alpha \Gamma &= -q\Gamma\alpha, & \delta \Gamma &= -q^{-1}\Gamma\delta + (q^{-1} - q)Ac, \\
bA &= qAb, & cA &= qAc, \\
bD &= q^{-1}Db + (q - q^{-1})\Omega\alpha, & cD &= q^{-1}Dc + (q - q^{-1})\Gamma\alpha, \\
b\Omega &= \Omega b, & c\Omega &= \Omega c + (q^2 - 1)A\alpha, \\
b\Gamma &= \Gamma b + (1 - q^2)A\alpha, & c\Gamma &= \Gamma c,
\end{aligned} \tag{25}$$

and

$$D_q u = q^2 u D_q, \quad u \in \{\alpha, b, c, \delta\}. \tag{26}$$

Before defining a coproduct on the algebra $\hat{\mathcal{A}}$, let us note the following facts. Let $\hat{T}$ and $\hat{T}'$ be any two supercommuting matrices that satisfy (1). We denote a product $\hat{T}\hat{T}'$ by T . Then, it can be verified that the matrix elements of T satisfy the commutation relations (6) of $GL_q(1|1)$, i.e. if

$$T = \begin{pmatrix} \alpha & b \\ c & \delta \end{pmatrix} \begin{pmatrix} \alpha' & b' \\ c' & \delta' \end{pmatrix}$$

then we have the relations (6). In short,

$$\hat{T}, \hat{T}' \in Gr_q(1|1) \implies T = \hat{T}\hat{T}' \in GL_q(1|1).$$

In view of these facts, we can say that there may be no coproduct of the usual form $\Delta(\hat{T}) = \hat{T} \dot{\otimes} \hat{T}$. For, this coproduct, if existed, would be invariant under the q -commutation relations (6) of $GL_q(1|1)$. But we can define a map on the algebra $\hat{\mathcal{A}}$ as follows:

$$\hat{\Delta} : \hat{\mathcal{A}} \longrightarrow \hat{\mathcal{A}} \otimes \hat{\mathcal{A}}, \quad \hat{\Delta}(\hat{T}) = \hat{T} \dot{\otimes} T + (-1)^{p(T)} T \dot{\otimes} \hat{T}. \tag{27}$$

Explicitly,

$$\begin{aligned}
\hat{\Delta}(\alpha) &= \alpha \otimes a + b \otimes \gamma + a \otimes \alpha - \beta \otimes c, \\
\hat{\Delta}(b) &= b \otimes d + \alpha \otimes \beta + a \otimes b - \beta \otimes \delta, \\
\hat{\Delta}(c) &= c \otimes a + \delta \otimes \gamma - \gamma \otimes \alpha + d \otimes c, \\
\hat{\Delta}(\delta) &= \delta \otimes d + c \otimes \beta - \gamma \otimes b + d \otimes \delta.
\end{aligned} \tag{28}$$

The action on the generators of $\hat{\mathcal{A}}$ of $\hat{\varepsilon} : \hat{\mathcal{A}} \longrightarrow \mathcal{C}$ is

$$\hat{\varepsilon}(\alpha) = \hat{\varepsilon}(b) = \hat{\varepsilon}(c) = \hat{\varepsilon}(\delta) = 0. \tag{29}$$

Finally, we define the coinverse as

$$\hat{S} : \hat{\mathcal{A}} \longrightarrow \hat{\mathcal{A}}, \quad \hat{S}(\hat{T}) = -(-1)^{p(T^{-1})} T^{-1} \hat{T} T^{-1}. \quad (30)$$

The action of $\hat{S}$ on the generators of $\hat{\mathcal{A}}$ is

$$\begin{aligned} \hat{S}(\alpha) &= -(\alpha A + b\Gamma)A + q(cA + \delta\Gamma)\Omega, \\ \hat{S}(b) &= -(\alpha A - c\Omega)\Omega - q(bA + \delta\Omega)D, \\ \hat{S}(c) &= -(\alpha A + b\Gamma)\Gamma - q(cA + \delta\Gamma)D, \\ \hat{S}(\delta) &= q^2(\alpha A - c\Omega)D + q(\alpha\Omega + q^2bD)\Gamma - (\alpha A + q^2\delta D)D. \end{aligned} \quad (31)$$

It is not difficult to check that the maps $\hat{\Delta}$ and $\hat{\varepsilon}$ are both algebra homomorphisms and $\hat{S}$ is an algebra anti-homomorphism and also the three maps satisfy the properties (16)-(18), and they preserve the relations (24) provided that the action on the generators of $\mathcal{A}$ of $\hat{\Delta}$ is the same with (13).

The coproduct, counit and coinverse which are specified above supply $\text{Gr}_q(1|1)$ with a structure, which can be called a quasi-Hopf algebra.

It is interesting to note that there is a close connection with the differential calculus⁶ on the quantum supergroup $\text{GL}_q(1|1)$ via the equation (23). In fact we have observed that the matrix elements of $\hat{T} \in \text{Gr}_q(1|1)$ are just the differentials of the matrix elements of $T \in \text{GL}_q(1|1)$. In other words, we can interpret the generating elements of $\text{Gr}_q(1|1)$ as differentials of coordinate functions on $\text{GL}_q(1|1)$. In this case, we can write $\hat{T} = dT$ (more information on these issues are given in ref. 6). Then the extended algebra can be interpreted as an algebra of differential forms on $\text{GL}_q(1|1)$. Thus the coproduct is interpreted as a (left and right) coaction of the quantum supergroup $\text{GL}_q(1|1)$ on differential forms. To this end, we consider the two maps

$$\Delta_R : \Gamma \longrightarrow \Gamma \otimes \mathcal{A}, \quad \Delta_R \circ d = (d \otimes \text{id}) \circ \Delta \quad (32a)$$

and

$$\Delta_L : \Gamma \longrightarrow \mathcal{A} \otimes \Gamma, \quad \Delta_L \circ d = (\tau \otimes d) \circ \Delta, \quad (32b)$$

where Γ denotes the differential algebra of $\mathcal{A}$. Here $\tau : \Gamma \longrightarrow \Gamma$ is the linear map of degree zero which gives $\tau(a) = (-1)^{p(a)}a$. We now define a map ϕ_R as follows

$$\phi_R(u_1 dv_1 + dv_2 u_2) = \Delta(u_1)\Delta_R(dv_1) + \Delta_R(dv_2)\Delta(u_2) \quad (33)$$

and another map ϕ_L by replacing L with R . The following identities are satisfied

$$(\phi_R \otimes \text{id}) \circ \phi_R = (\text{id} \otimes \Delta) \circ \phi_R \quad (\text{id} \otimes \epsilon) \circ \phi_R = \text{id}, \quad (34a)$$

and

$$(\text{id} \otimes \phi_L) \circ \phi_L = (\Delta \otimes \text{id}) \circ \phi_L \quad (\epsilon \otimes \text{id}) \circ \phi_L = \text{id}. \quad (34b)$$

Consequently, we define the map $\hat{\Delta}$, in (27), as

$$\hat{\Delta} = \phi_R + \phi_L. \quad (35)$$

ACKNOWLEDGEMENT

This work was supported in part by **T. B. T. A. K.** the Turkish Scientific and Technical Research Council.

I would like to express my deep gratitude to the referee for critical comments and suggestions on the manuscript.

- ¹ N. Y. Reshetikhin, L. A. Takhtajan and L. D. Faddeev, Leningrad Math. J. **1** (1990), 193.
- ² V. G. Drinfeld, Quantum groups, in *Proc. IMS, Berkeley*, (1986).
- ³ Yu. I. Manin, Commun. Math. Phys. **123** (1989), 163;
E. Corrigan, D. Fairlie, P. Fletcher and R. Sasaki, J. Math. Phys. **31** (1990), 776;
S. Schwenk, B. Schmidke and S. Vokos, Z. Phys. C **46** (1990), 643;
B. Schmidke, S. Vokos, and B. Zumino, Z. Phys. C **48** (1990), 249.
- ⁴ S. Çelik and S. A. Çelik, Balkan Phys. Lett. **3** (1995), 188.
- ⁵ S. Çelik, J. Math. Phys. **37** (1996), 3568;
S. Çelik, Balkan Phys. Lett. **5** (1997), 149.
- ⁶ S. Çelik and S. A. Çelik, "On the differential geometry of $GL_q(1|1)$ ", J. Phys. A: Math. Gen. (in press) (1998).